### polar. mtg. 07.03.12

# pC polarimeter data


- Result from 1 chan. MUX bypass
- Brief look at waveforms (Andrei)
- Data cleaning ⇒ improving polarization measurement

#### Extras:

- Special target data recorded so far
- History: demise of Y2U det. 4 preamp

MUX bypass test

Access 29.02.12 test of "MUX reflection" hypothesis:
 Y1D ch. 25 output bypassed around MUX, direct to DAQ
 Y2U ch. 25 left unconnected




MUX bypass did not remedy this particular problem

# Waveform readout


Thanks to Andrei: adapted his tool for AGS→RHIC data, and set up for me to use

- Can readout and display RAW sample data from WFDs
- Waveforms recorded for triggered events: max. sample > threshold, within T-window (I think...)
- Samples digitized every 2.4 nS (= 2 usual TDC units)
- Data taken during development period 01.03.12
- After Haixin took usual carbon sweep, I took no-target run w/ samples
- Here data from Y2U with run 16495.301:
  - 200 MHz V=20kV
  - 12 filled bunches equally spaced, except bcn-60 empty




- A quiet (no noise) chan. 13 (det. 2): no target, only pulser events
- Main pulse every 180 rev. + 1 bunch; 'echo' pulse 86 bunches later


### Run=16495.0 Det=1 Strip=12



• Ch. 13 superposition all event waveforms:




• This chan. in pC sweep run:




Note apparent T difference:

- Waveform peak ~75
- on banana plot T~68
- CFD: "const. frac. discrim." extrapolate back from peak to ~1/3 max. pulse height


- Next noisy chan. 36 (det. 3): triggered every filled bunch
- Note bunch 70 (after long gap) different
- 1st 12 events: Run=16495.0 Det=2 Strip=35




- Next noisy chan. 36 (det. 3): triggered every filled bunch
- Note bunch 70 (after long gap) different
- 2<sup>nd</sup> 12 events: Run=16495.0 Det=2 Strip=35




Ch. 36 superposition all event waveforms:
 This chan. in pC sweep run:




- The 2<sup>nd</sup> blob on TDC-Amp plot is bcn 70 after long gap:
- Wider pulse: wave peak 75-80 const. frac. discrim. ~55





- Stranger: chan. 2 (det. 1): triggered every filled bunch
- Prompt pulse visible near T=10;
- After-pulse extending to next bunch, also triggered

Run=16495.0 Det=0 Strip=1



# Data cleaning

- Effects seen in recorded waveforms already seen on scope
- But also: they reflect the nature of recorded data:
  noise pulse are wider (T) than signals (carbon pulses ~ pulser)

#### WFD reconstructed pulse info:

- Amp = amplitude max. sample
- Tdc = CFD time, resolution 1.2 nS
- Int = integral (sum) of all samples
- TdcAMax = time of max. sample, resolution 2.4 nS


#### Pulse T-width related to:

- Int/Amp ~ width of pulse
- TdcAMax-Tdc ~ length of pulse rising edge

### Should be able to use (Int/Amp), (TdcAMax-Tdc) to identify noise


Look at pC sweep data same run 16495.301 \( \square\)

# Int/Amp



- Noise pulses have larger Int/Amp
- Pretty clean signal/noise separation

## TdcAMax-Tdc



- Noise pulses have larger TdcAMax-Tdc
- Pretty clean signal/noise separation

# Data cleaning

Clearly can use tools like these to clean our data:

- Identify/reject noise events
- Identify/reject noisy channels
- Identify/reject noisy detectors

Already simple versions of channel cleaning are in use and giving respectable polarization measurements:

- Offline (for over 1 week): cut on topography of Tdc-Amp plane
- Online (since yesterday): reject detectors large variation #events/chan. (new online values not archived, sorry Phil...)
- Not the whole story: next need to worry about baseline (basecurve?)
  shifts under good carbon signal ⇒ shifts of E-window for pol. measure
- But we definitely will extract good polarization measurements with this data as is

# Data cleaning

Clearly can use tools like these to clean our data:

- Identify/reject noise events
- Identify/reject noisy channels
- Identify/reject noisy detectors

Already simple versions of channel cleaning are in use and giving respectable polarization measurements:

- Offline (for over 1 week): cut on topography of Tdc-Amp plane
- Online (since yesterday): reject detectors large variation #events/chan. (new online values not archived, sorry Phil...)
- Not the whole story: next need to worry about baseline (basecurve?)
  shifts under good carbon signal ⇒ shifts of E-window for pol. measure
- But we definitely will extract good polarization measurements with this data as is

### **Very important:**

- Need stable running configuration, channels on and read out
- Must avoid random ill-motivated perturbations on system

# Extras

# Special target data

Data recorded with special targets (from data base)

### Highly twisted targets

- B1U H3 0 runs
- B1U H6 0 runs
- B2D V3 40-50 runs Feb. 13-14
- B2D V6 0 runs

#### **Short targets**

- B1U H5 0 runs
- B2D V5 0 runs

#### Wide target

B2D H4 – 10 runs Feb. 28-29

### Very wide target

B2D H6 – 2 runs March 1

Special targets (nominal dimensions except as noted) and locations are:

- 4 highly twisted targets (B1U H3,H6; B2D V3,V6)
- 3 short targets 20 mm long (B1U H5; B2D V5; Y1D H6)
- 1 wide target 50-75 μ wide (B2D H4)
- 1 very wide target 5 mm wide, fixed angle w.r.t. beam, detectors (B2D H6)

# Demise of Y2U det. 4 preamp

#### 20.2 ~11:55

Last OK run 16449.307; usual 'noise' effect, otherwise normal


←nothing notable in e-log here, quick refill

#### 20.2 ~13:30

- Run 16450.302 start problems, 2 chan. weak or dead or messy
- Continued throughout fill 16450

#### 21.2 ~02:30

- Run 16451.301 start disaster, all chan. ~dead, plot empty
- Run 16451.303 start crazy behavior:



# Int/Amp vs. TdcAMax-Tdc

Contours: good chans. det. 2

Boxes: noisy chan. 37

